

The Cyclic Dinucleotide c-di-AMP Is an Allosteric Regulator of Metabolic Enzyme Function

Kamakshi Sureka,^{1,8} Philip H. Choi,^{2,8} Mimi Precit,¹ Matthieu Delince,^{3,9} Daniel A. Pensinger,⁴ TuAnh Ngoc Huynh,¹ Ashley R. Jurado,² Young Ah Goo,⁵ Martin Sadilek,⁶ Anthony T. Iavarone,⁷ John-Demian Sauer,⁴ Liang Tong,^{2,*} and Joshua J. Woodward^{1,*}

¹Department of Microbiology, University of Washington, Seattle, WA 98195, USA

²Department of Biological Sciences, Columbia University, New York, NY 10027, USA

³Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA

⁴Department of Medical Microbiology and Immunology, University of Wisconsin-Madison, Madison, WI 53706, USA

⁵Department of Pharmaceutical Sciences, University of Maryland, Baltimore, MD 21201, USA

⁶Department of Chemistry, University of Washington, Seattle, WA 98195, USA

⁷QB3/Chemistry Mass Spectrometry Facility, University of California, Berkeley, CA 94720, USA

⁸Co-first author

⁹Present address: School of Life Sciences, Swiss Federal Institute of Technology in Lausanne (EPFL), Lausanne 1015, Switzerland

*Correspondence: ltong@columbia.edu (L.T.), jjwoodwa@uw.edu (J.J.W.)

<http://dx.doi.org/10.1016/j.cell.2014.07.046>

SUMMARY

Cyclic di-adenosine monophosphate (c-di-AMP) is a broadly conserved second messenger required for bacterial growth and infection. However, the molecular mechanisms of c-di-AMP signaling are still poorly understood. Using a chemical proteomics screen for c-di-AMP-interacting proteins in the pathogen *Listeria monocytogenes*, we identified several broadly conserved protein receptors, including the central metabolic enzyme pyruvate carboxylase (LmPC). Biochemical and crystallographic studies of the LmPC-c-di-AMP interaction revealed a previously unrecognized allosteric regulatory site 25 Å from the active site. Mutations in this site disrupted c-di-AMP binding and affected catalytic activity of LmPC as well as PC from pathogenic *Enterococcus faecalis*. C-di-AMP depletion resulted in altered metabolic activity in *L. monocytogenes*. Correction of this metabolic imbalance rescued bacterial growth, reduced bacterial lysis, and resulted in enhanced bacterial burdens during infection. These findings greatly expand the c-di-AMP signaling repertoire and reveal a central metabolic regulatory role for a cyclic dinucleotide.

INTRODUCTION

The colonization of diverse environmental and host niches by bacteria depends on the ability to sense and respond to external stimuli. Signal transduction pathways dependent upon nucleotide-based second messengers are ubiquitous

among bacteria (Kalia et al., 2013) and mediate cellular development, group behavior, and metabolism as a result of changing environmental conditions. Among such molecules, cyclic di-adenosine monophosphate (c-di-AMP, cdA) has recently emerged as a broadly conserved second messenger of fundamental importance for microbial growth and physiology, with effects on stress responses, antibiotic resistance, cellular morphology, bacterial growth, and virulence (Campos et al., 2014; Corrigan et al., 2011; Corrigan et al., 2013; Luo and Hellmann, 2012; Mehne et al., 2013; Oppenheimer-Shaanan et al., 2011; Pozzi et al., 2012; Smith et al., 2012; Witte et al., 2013; Zhang and He, 2013). Moreover, c-di-AMP secreted by *L. monocytogenes*, *M. tuberculosis*, and *C. trachomatis* during infection results in an IFN- β -mediated host immune response (Barker et al., 2013; Woodward et al., 2010; Yang et al., 2014). Together, these reports have established c-di-AMP as a crucial intracellular signaling molecule and an important component of innate immune detection during infection. Despite this central importance, the molecular mechanisms of c-di-AMP-mediated signal transduction in bacteria are just being elucidated.

Components of second messenger signaling systems include mechanisms to generate, degrade, and relay the signal. C-di-AMP is synthesized from two molecules of ATP (Witte et al., 2008), and in some cases ADP (Bai et al., 2012), through the catalytic activity of di-adenylate cyclases (DACs). The signal is subsequently degraded to pApA or AMP by DHH/DHHA1 domain containing phosphodiesterases (Bai et al., 2013; Rao et al., 2010). Second messengers initiate signal transduction by binding to and altering protein and nucleic acid function, resulting in transcriptional, translational, and posttranslational alterations within a cell. Recent reports have identified several mechanisms of c-di-AMP regulation within these contexts, including riboswitches and protein receptors




Figure 1. Identification of c-di-AMP-Interacting Proteins from *L. monocytogenes*

(A) Schematic diagram of chemical proteomics used to identify c-di-AMP-interacting proteins. 1° and 2° represent direct and indirect specific binding proteins.

(B) Representative pull-down from bacterial lysates with c-di-AMP (+) and control sepharose (-). The remaining lanes from the SDS-PAGE gel have been removed for clarity.

(C) Quantitative shotgun proteomics of c-di-AMP-binding proteins. Data are sorted based upon fold change (FC) in spectral count ratio (c-di-AMP sepharose/control sepharose). Data points in red represent proteins with FC > 7 (horizontal black line) and a false discovery rate (FDR) < 0.05.

(D) List of top hits identified by chemical proteomics. Studies performed in *L. monocytogenes* strain 10403S. Locus numbers based on strain EGD-e (Accession number: NC_003210.1).

(E and F) Proteins in bold confirmed for direct c-di-AMP binding with (E) purified recombinant proteins or (F) crude cell lysate for proteins identified by chemical proteomics.

Data are mean \pm SEM and are representative of multiple independent experiments. * p < 0.05 (Students *t* test, two-tailed). See also Figure S1, S2, and S3.

(Bai et al., 2014; Corrigan et al., 2013; Nelson et al., 2013; Zhang et al., 2013). However, c-di-AMP riboswitches are absent in some organisms and protein receptor homologs are not well conserved. Similarly, the functional roles of identified protein receptors cannot explain the breadth of phenotypes associated with altered c-di-AMP levels. Together these observations imply that additional c-di-AMP signaling pathways likely exist.

Listeria monocytogenes is a gram-positive intracellular bacterial pathogen that provides an attractive model to interrogate the molecular details of protein-mediated c-di-AMP signaling. C-di-AMP produced from a single DAC containing protein is a central regulator of numerous fundamental processes within this organism, including growth, cell wall homeostasis, stress responses, and the establishment of infection (Witte et al., 2013). In this study we developed a chemical proteomics approach to identify the c-di-AMP-interacting proteins. We identified several c-di-AMP-binding proteins that are broadly conserved in numerous bacterial species. Among these proteins, we structurally and biochemically characterized the interaction with pyruvate carboxylase (PC) and characterized the consequences of c-di-AMP regulation of central metabolic activity on *L. monocytogenes* survival within the host environment. Our findings provide insight into the central biological function of c-di-AMP among the diverse bacteria known to generate this second messenger.

RESULTS

Chemical Proteomics to Define the c-di-AMP Interactome of *L. monocytogenes*

We utilized a proteomics approach to identify the c-di-AMP interactome of *L. monocytogenes* (Figure 1A). The nucleotide was covalently conjugated to epoxy-activated sepharose beads (Figure S1). Nucleotide-bound and control beads were used in parallel for affinity purification of proteins from *L. monocytogenes* 10403S. Initial binding results were visually analyzed by SDS-PAGE (Figure 1B) and subsequently using gel-free quantitative shotgun proteomics (Figure 1C). We identified 12 proteins that were statistically significant among replicates (false discovery rate < 0.05) and enriched > 7-fold by spectral count ratio (>2 SD) with c-di-AMP versus control sepharose (Figure 1D). The four proteins with highest enrichment are of conserved hypothetical annotation, with homologs found in a large number of bacteria. Among these, the homolog of Lmo2692 in *S. aureus* was recently identified as a c-di-AMP-binding protein (Corrigan et al., 2013). In addition, the known c-di-AMP phosphodiesterase PdeA (Lmo0052) is the fifth protein on our list (Witte et al., 2013). The successful identification of known c-di-AMP-binding proteins validated our approach.

To confirm these interactions, proteins were probed for direct c-di-AMP-binding by differential radial capillary action of ligand




Figure 2. LmPC Specifically Binds and Is Inhibited by c-di-AMP.

(A) Binding titration of c-di-AMP and LmPC using ^{32}P -c-di-AMP.

(B) Enzymatic activity of LmPC in the presence of Acetyl-CoA (100 μM), varied pyruvate, and c-di-AMP concentrations as indicated.

(C) Rate constants of pyruvate turnover in the presence of c-di-AMP.

(D) Enzymatic activity of LmPC in the presence of pyruvate (20 mM), varied Acetyl-CoA (allosteric activator) and c-di-AMP concentrations as indicated.

(E) LmPC activity in the presence of nucleotide analogs. Data reported as percent activity relative to that without added nucleotide. All data reported as the mean \pm SEM and are representative of multiple independent experiments. See also Figure S4.

assay (DRaCALA) (Roelofs et al., 2011). ^{32}P -labeled c-di-AMP was synthesized (Figure S2) and used to quantify binding. Those proteins predicted to be soluble by in silico analysis (Bernsel et al., 2009) were expressed and purified to homogeneity from *E. coli*. The three soluble proteins of conserved hypothetical annotation (renamed here PstA, CbpA, and CbpB) as well as PdeA and LmPC exhibited binding in purified form (Figure 1E). No binding was observed for purified recombinant ClpC, OppA, CtaP, or Lmo1634, which were likely identified because they interact with a direct c-di-AMP-interacting protein (labeled 2° in Figure 1A) or contain posttranslational modifications that facilitate c-di-AMP binding. NrdR exhibited significant instability in purified form, while Lmo1466 and NADH dehydrogenase are predicted to be membrane anchored. As such, these proteins were characterized in whole-cell lysates. *E. coli* lysates containing either an empty vector or encoding Lmo0553 were used as negative and positive controls, respectively (Figure 1F). A significant level of binding was observed for Lmo1466 and NrdR-expressing lysates, consistent with a direct interaction with c-di-AMP. No interaction was observed for lysates derived from *E. coli* expressing NADH dehydrogenase. However, overexpression of NADH dehydrogenase was not observed and the direct interaction with c-di-AMP was therefore inconclusive.

These observations support a specific interaction with c-di-AMP.

c-di-AMP is an Allosteric Inhibitor of LmPC

Among those newly identified c-di-AMP-binding proteins, only LmPC (120 kDa) has a known biological function. In fact, LmPC is essential for *L. monocytogenes* growth (Schär et al., 2010). Therefore, we focused our initial studies on the detailed characterization of this molecular interaction and its biological consequences. PC is a multidomain enzyme that catalyzes the ATP-dependent carboxylation of pyruvate to oxaloacetate. During catalysis, the biotin carboxyl carrier protein (BCCP) domain is carboxylated in an ATP-dependent manner by the biotin carboxylase (BC) domain. Subsequently, the carboxyltransferase (CT) domain mediates transfer of the carboxyl from biotin to the substrate pyruvate. The PC tetramerization (PT) domain is important for the quaternary structure of the enzyme (St Maurice et al., 2007; Xiang and Tong, 2008) and together with the BC domain defines the binding site of the allosteric activator acetyl-CoA (St Maurice et al., 2007; Yu et al., 2009).

Binding studies indicated that LmPC interacts with c-di-AMP with a K_d of 8 ± 0.2 μM (Figure 2A). We observed diminished PC activity in the presence of increasing concentrations of c-di-AMP ($K_i \sim 3$ μM , Figure 2B). The inhibitory effect was at the




Figure 3. Crystal Structure of LmPC in Complex with c-di-AMP

(A) Omit F_o-F_c electron density map for c-di-AMP at 2.5 Å resolution, contoured at 3σ .

(B) Schematic drawing of the structure of LmPC tetramer in complex with two c-di-AMP molecules. The domains of monomer 1 are colored separately, according to the diagram at the bottom of the panel. c-di-AMP is shown as a sphere model and labeled cdA (carbon atoms in black). The metal ion in the active site of the CT domain is shown as a gray sphere and labeled M^{2+} . The distance from the CT active site and c-di-AMP-binding site is indicated with the dashed red line.

(C) Detailed interactions between c-di-AMP and LmPC. Hydrogen-bonding interactions are indicated with dashed lines (in red). Water molecules are shown as red spheres. All structure figures were produced with PyMOL (www.pymol.org). See also Table S1 and Figure S5.

level of the apparent k_{cat} with slight lowering of the apparent K_m (Figure 2C), indicating that c-di-AMP does not compete with the pyruvate substrate. We also found that the inhibition by c-di-AMP could be overcome by high concentrations of acetyl-CoA (Figure 2D). Finally, the inhibition was specific to c-di-AMP as most of structurally related dinucleotides (Figure S4), including c-di-GMP, bacterial cGAMP (cyclic[Gp(3',5')Ap(3',5')]), or pApA, the breakdown product of c-di-AMP, showed no effect on LmPC activity, while c-di-IMP did show weak inhibitory activity (Figure 2E). Overall, these biochemical studies define c-di-AMP as an allosteric inhibitor of LmPC.

Binding Mode of c-di-AMP in LmPC

To understand the molecular basis for the enzyme regulation, we determined the structure of the LmPC-c-di-AMP complex at 2.5 Å resolution (Table S1). Clear electron density was observed for the c-di-AMP ligand based on the crystallographic analysis (Figure 3A). Unexpectedly, we observed another c-di-AMP that bridges two c-di-AMP molecules bound to neighboring LmPC in this crystal (Figure S5A). The bridging c-di-AMP inserts one of its adenine bases between the two bases of the bound c-di-AMP, forming π -stacking interactions, but it has no direct contacts with the protein (Figure S5A). We observed a second crystal form of the complex from the same crystallization condi-

tion, and determined its structure at 2.7 Å resolution (Table S1). The bridging c-di-AMP molecule is absent in this crystal (Figure S5B), but the structure of the complex is essentially identical to the other crystal form (Figure S5C). Therefore, the bridging c-di-AMP is most likely an artifact in that crystal form and will not be discussed further. A citrate molecule was found in the acetyl-CoA-binding site, although our kinetic studies showed that citrate had only small effects on the catalytic activity of LmPC. Its presence in the structure was likely due to the high concentration in the crystallization solution.

LmPC is a tetramer in the crystal structure (Figures 3B and S5C), consistent with solution studies as well as earlier structures of *R. etli* PC (RePC), *S. aureus* (SaPC), and *H. sapiens* (HsPC) (St Maurice et al., 2007; Xiang and Tong, 2008). The tetramer is composed of two layers, with two monomers in each layer. There are extensive interactions between the two layers, mediated through the BC and CT domain dimers, which are located at opposing corners of the diamond-shaped tetramer. In contrast, there are few direct contacts between the two monomers in the same layer. All four BCCP domains are observed, but the linkers between the PT and the BCCP domains (residues 1060–1066) are disordered. As a result, each BCCP domain is assigned to a monomer based on the earlier structure of SaPC (Figure 3B), although the actual PT-BCCP

connectivity is ambiguous in LmPC. The overall position of BCCP is similar to that in the exo site of the SaPC structure (Xiang and Tong, 2008), but its orientation is very different (Figure S6A). Consequently, the biotin group in the current structure is projected into the solvent and disordered. Besides the PT-BCCP linker, the B domain of BC (residues 135–205) is also disordered in the crystals.

The c-di-AMP nucleotide is bound at the dimer interface of two CT domains (Figures 3B), with the 2-fold symmetry axis of the compound aligned with that of the CT domains (Figure 3C). Therefore, each LmPC tetramer binds only two c-di-AMP molecules (Figure 3B). The binding site is formed by two helices in the TIM barrel fold of the CT domain (Figure 3C), but it is located ~25 Å away and on the opposite face from the CT active site (Figures 3B). This region of PC has not been known previously to have any regulatory or structural significance.

The c-di-AMP ligand adopts a U-shaped conformation (Figure 3A) and interacts with LmPC through hydrophobic and hydrogen-bonding interactions (Figure 3C). The aromatic ring of Tyr722 has a direct face-to-face π -stacking arrangement with an adenine base. One of the terminal oxygen atoms of the phosphate group of c-di-AMP forms a hydrogen bond with the hydroxyl group of Tyr749, while the other terminal oxygen interacts with the Ser756 side chain through a water molecule. The ribose is located near Ala753, which together with Ala752 may provide the space necessary for the ligand to bind.

The U-shaped conformation of c-di-AMP is similar to the binding mode of c-di-GMP in some proteins, such as the bacterial PilZ domain (Benach et al., 2007) and the human STING protein (Huang et al., 2012; Ouyang et al., 2012; Shang et al., 2012; Shu et al., 2012; Yin et al., 2012). In addition, c-di-GMP is also aligned with the 2-fold symmetry axis of a STING dimer, and the guanine base is π -stacked with a Tyr side chain. Together these observations identify a binding site of c-di-AMP to LmPC and confirm kinetic studies that the site is unique from known substrates or allosteric regulators of this family of enzymes.

The c-di-AMP-Binding Site Affects PC Catalytic Activity

To assess the functional importance of the binding site, we introduced amino acid mutations into LmPC. We noted conservation of a π -stacking aromatic amino acid among many cyclic dinucleotide-binding proteins, including STING, VpsT, and the EAL domain protein YahA (Huang et al., 2012; Krasteva et al., 2010; Ouyang et al., 2012; Shang et al., 2012; Sundriyal et al., 2014; Yin et al., 2012), implicating Y722 as a critical residue for nucleotide recognition. Two amino acid substitutions (Y722F and Y722T) were introduced into this site. The Y722F mutant was chosen to assess the importance of the hydroxyl group interacting with the N6 of the adenine base. The Y722T mutant was chosen due to the presence of this amino acid in the corresponding site of HsPC (Figure 4A). Comparison of the LmPC and SaPC structures revealed that the analogous binding pocket does not apparently exist within the latter enzyme (Figures 4B, 4C, S6A, and S6B). This can be attributed to the presence of a Lys and Gln substitution in SaPC corresponding to A752 and A753 of LmPC, respectively, which introduce significant steric bulk into the pocket. Similarly, the Y749 residue required for hydrogen bonding to the terminal oxygen of the phosphate at the bottom of

this pocket has been lost. These observations were used to guide amino acid substitutions in the LmPC protein.

The mutants were expressed and purified following the same protocol as was used for the wild-type protein. All amino acid mutants, except Y749L, were catalytically active. However, the Y722T, A752K, and A753Q variants lost sensitivity to inhibition by c-di-AMP even at 100 μ M concentration (Figure 4D). Interestingly, the enzyme activity in the absence of c-di-AMP was substantially different from WT, with both increased and reduced activity being observed for the mutants, suggesting that this region of the enzyme has a functionally relevant effect on PC catalysis. The Y722F mutant behaved similar to the wild-type enzyme in terms of its catalytic activity and responsiveness to c-di-AMP, establishing that the hydroxyl group of Y722 is dispensable for nucleotide recognition. In agreement with the kinetic data, binding to c-di-AMP was only observed for Y722F and none of the other amino acid variants (Figure 4E).

We noted that although several of the residues required for c-di-AMP binding by LmPC are absent in other bacterial enzymes, the required chemical features of the binding pocket might be conserved. Specifically, when comparing LmPC to PC from lactic acid bacteria (Figure 4A), such as PC from the opportunistic pathogen *Enterococcus faecalis* (EfPC), structural modeling predicted that the A752S and S756K variants could establish favorable interactions with the c-di-AMP phosphate group and compensate for the loss of the Y749 residue (Figures 4F–4G). Indeed, recombinant EfPC binds c-di-AMP and its catalytic activity is inhibited by the nucleotide (Figures 4G and 4H). Consistent with a conserved binding site between the enzymes, mutation of the base stacking tyrosine results in loss of c-di-AMP binding and regulation. Similarly, SaPC is insensitive to c-di-AMP. Overall, the results from these studies confirm the c-di-AMP-binding site identified by the structural analysis and identify a previously unrecognized region important for PC regulation in *L. monocytogenes* and other bacteria.

C-di-AMP Binding Induces Large Conformational Changes in LmPC

We have also determined the structure of apo LmPC (Table S1), in the absence of c-di-AMP. The best quality X-ray diffraction data set we could collect after screening through a large number of crystals was at 3.3 Å resolution. There are two tetramers in the asymmetric unit that have essentially the same conformation, with root mean square (rms) distance of 0.1 Å among their equivalent $C\alpha$ atoms (NCS restraints were applied throughout the refinement and the crystal had pseudo C2 space-group symmetry). Therefore, only one of these tetramers will be used for further analysis. All four BCCP domains and residues 850–950 in the CT domain of one of the monomers had poor electron density and were not modeled. The B domain of BC was partially ordered in the four monomers and there was weak density for an ATP molecule.

A comparison of the structure of apo LmPC and that bound to c-di-AMP showed dramatic rearrangements in the tetramer upon ligand binding (Figure 5A). Near the c-di-AMP-binding site, there are significant changes in the orientation and position of the two CT domains of the dimer, corresponding to a rotation of ~4.5° for each CT domain (Figure 5B). As a consequence, the

A

<i>L. monocytogenes</i>	719	QAAYRL	724	747	GIYTYAAAVSAG	758
<i>S. aureus</i>		KAAYEL			GLLTYKQAI DAG	
<i>H. sapiens</i>		TACTML			GVAAMLACAQAG	
<i>E. faecalis</i>		QAAYML			GIITYSAATKAG	
<i>L. farvieae</i>		QAAYML			GIITYSGATKAG	
<i>L. lactis</i>		QAAYML			GIITYSGAICAG	

Figure 4. Functional Analysis of the c-di-AMP-Binding Site of PC

(A) Conservation of residues in the c-di-AMP-binding site. The five residues in the binding site are highlighted in red.

(B) Molecular surface of the LmPC-binding site for c-di-AMP (labeled cdA), colored by the electrostatic potential (red: negative, blue: positive).

(C) Molecular surface of SaPC at the CT dimer interface. The c-di-AMP molecule is shown as a reference in stick models.

(D) Effects of c-di-AMP on the catalytic activity of wild-type and mutant LmPC.

(E) C-di-AMP binding to WT and mutant LmPC proteins measured by DRaCALA.


(F) LmPC c-di-AMP-binding pocket with corresponding EfPC amino acid variations.

(G) Surface view of modeled EfPC.

(H) C-di-AMP binding to WT and Y718T EfPC.

(I) Activity of WT and Y718T EfPC in the presence and absence of c-di-AMP.

Data are mean ± SEM (D, E, G, and H) and are representative a multiple independent experiments. See also Figure S6.






Figure 5. Large Conformational Differences in the Structure of apo LmPC

(A) Overlay of the structure of LmPC in complex with c-di-AMP (in color) with that of apo LmPC (in gray). The superposition was based on monomer 1. Large differences in the positions of the other monomers are visible.

(B) Conformational differences in the c-di-AMP-binding site in the structure of apo LmPC.

(C) Molecular surface of apo LmPC at the CT dimer interface. The c-di-AMP molecule would clash with the enzyme in this conformation.

(D) Overlay of the four monomers of the LmPC tetramer in the complex with c-di-AMP. The superposition is based on the CT domain only. A small conformational difference is seen for the BC domain.

(E) Overlay of the four monomers of the apo LmPC tetramer. Large differences are observed in the positions and orientations of the BC domains.

domains are overlaid for the apo enzyme, and the hinge rotation is located near the boundary between the BC and PT domains (Figure 5E). In comparison, the overlay of the CT domains brought the PT domains into superposition. The PT domains of SaPC and HsPC have important roles in mediating contacts in the tetramer (Xiang and Tong, 2008). The PT domains in the c-di-AMP complex of LmPC and apo LmPC are positioned further away from each other and do not make direct contacts (Figure 5A).

Metabolic Balance Mediated by c-di-AMP Affects Intracellular Growth of *L. monocytogenes*

Structural and biochemical characterization of the interaction between LmPC and c-di-AMP implicates the nucleotide as a metabolic regulator in

side chains of both Tyr722 and Ser756 in the apo enzyme would clash with the c-di-AMP molecule in the complex. These changes greatly reduce the size of the pocket in this region, such that it can no longer accommodate c-di-AMP (Figure 5C).

The BC dimer in apo LmPC has roughly the same structure as that in the complex with c-di-AMP (Figure S5D). However, the position and orientation of the BC domain relative to the CT domain are different between the apo enzyme and the c-di-AMP complex. In addition, the four LmPC monomers in the c-di-AMP complex have essentially the same conformation, while the monomers in the apo enzyme have large structural differences. This is best visualized by overlaying the CT domains of the four monomers in each structure. For the c-di-AMP complex, overlaying the CT domains essentially brought the BC and PT domains into overlay as well, with a difference of only $\sim 1^\circ$ in their orientations (Figure 5D). In comparison, the four monomers differ by $\sim 10^\circ$ in the orientations of their BC domains after the CT

L. monocytogenes. *Listeria* species have an incomplete citric acid cycle, and the oxaloacetate (OA) generated by LmPC is used as a precursor for aspartate and glutamate biosynthesis (Figure 6A) (Kim et al., 2006). To study the effects of c-di-AMP levels on metabolism in *L. monocytogenes*, we measured incorporation of isotope label from U- ^{13}C -glucose into amino acids alanine (Ala), aspartate (Asp), and glutamate/glutamine (Glx) in both WT and a conditional depletion strain of DacA (*cΔdacA*), which exhibits diminished levels of c-di-AMP (Witte et al., 2013). Bacteria with depleted c-di-AMP exhibited substantially increased ^{13}C incorporation into Glx relative to bacteria with normal levels of the nucleotide (Figure 6B). De novo synthesis of Glx is entirely dependent on LmPC activity in *L. monocytogenes* (Schär et al., 2010) and flux through the oxidative branch of the citric acid cycle. To determine if these observations are a result of PC activity directly, we generated a double conditional depletion strain, *cΔdacA cΔpycA*, with IPTG inducible expression of




Figure 6. Metabolic Imbalance Specifically Alters *L. monocytogenes* Intracellular Growth

(A) De novo biosynthesis of amino acids and labeling patterns from ^{13}C -glucose by *L. monocytogenes*. Dark spheres, ^{13}C . Light spheres, natural abundance carbon.

(B) ^{13}C enrichment into Ala, Asp, and Glx from midexponential *L. monocytogenes* strains.

(C) Magnitude of growth of *L. monocytogenes* strains in iBMMs. Data reported as the ratio of CFU recovered at 8 hpi relative to 0.5 hpi.

(D) ^{13}C enrichment into Ala, Asp, and Glx from midexponential *L. monocytogenes* strains.

(E) Immortalized murine bone-marrow-derived macrophages were infected with *L. monocytogenes* and CFU were enumerated at the indicated times.

(F) Plaque area from mouse fibroblasts (L2 cells) infected with indicated strains and normalized to WT.

(G) Bacterial burden in organs 48 hr postinfection with the median indicated by a horizontal bar.

(C–F) Data are mean \pm SEM (B–G) Data are representative of at least two independent experiments. (D) * $p < 0.001$ (Students t test, two-tailed). See also Figure S7.

DacA and theophylline inducible expression of PC (Figure S7A). Indeed, increased levels of theophylline in the absence of IPTG led to a direct increase in Glx synthesis (Figure 6B). Thus, c-di-AMP, through its action on LmPC and possibly other downstream targets that affect flux through this pathway, negatively affects de novo Glx biosynthesis. In comparison, reduced c-di-AMP levels did not affect Asp synthesis, indicating that LmPC is not the limiting activity for this pathway.

We hypothesized that this metabolic imbalance may be responsible for the defects in bacterial growth and host colonization associated with low levels of c-di-AMP. To test this possibility we infected macrophages with the *c* Δ *dacA* *c* Δ *pycA* strain grown in the presence of varying concentrations of theophylline. Increased cytosolic growth was observed as theophylline levels were increased until a critical threshold was achieved, after which the strain exhibited limited growth comparable to *c* Δ *dacA* (Figure 6C). These observations are consistent with previous reports where low PC activity results in diminished bacterial growth in the cytosol. However, our findings also show that enhanced PC activity is detrimental in the absence of c-di-AMP and are consistent with the requirement for balanced metabolic activity to establish infection in the host.

It has not been established why PC is essential for *L. monocytogenes*, though disruption of *aspB* resulted in diminished bacterial growth (data not shown), consistent with this being the essential branch emanating from the precursor metabolite oxaloacetate. Based on these observations, we hypothesized that the metabolic imbalance associated with diminished

c-di-AMP levels was a consequence of elevated flux to Glx. To segregate the detrimental effects of hypo and hyperactive PC we deleted citrate synthase (CitZ, Figure 6A), the first committed step to Glx biosynthesis. Incorporation of ^{13}C into Glx was abolished in the WT and *c* Δ *dacA* strains containing the *citZ* mutation (Figure 6D), while levels of Asp and Ala incorporation remained unchanged. Thus, disruption of *citZ* prevents unregulated glutamate/glutamine biosynthesis from glucose under conditions of low c-di-AMP.

Having corrected the elevated synthesis of Glx, we detailed the effects of metabolic imbalance associated with altered c-di-AMP signaling on *L. monocytogenes* growth and infection. Although disruption of *citZ* had no discernable effect on broth growth in the presence or absence of c-di-AMP production (Figure S7B), a significant increase in intracellular growth in the absence of c-di-AMP was observed within immortalized bone-marrow-derived macrophages (iBMM) and mouse fibroblasts (Figures 6E and 6F). No change in growth was observed for *citZ* disruption in the WT background (Figures 6F and S7C). Analysis of bacterial growth revealed that the doubling time was not significantly altered by loss of *citZ*, consistent with observations in rich media. However, the ability of *L. monocytogenes* to maintain intracellular growth between 5 and 8 hr was rescued by disruption of *citZ*, resulting in nearly WT levels of bacterial burden at later time points. Expression of the citrate synthase gene *in trans* attenuated intracellular growth of the *c* Δ *dacA* Δ *citZ* strain back to *c* Δ *dacA* levels (Figure 6E).




Figure 7. C-di-AMP-Induced Metabolic Imbalance Provokes Intracellular Bacteriolysis and Host Cell Pyroptosis

(A) Cytotoxicity induced in iBMMs by the indicated strains following infection was measured by LDH release.

(B) Intracellular lysis of bacterial strains in iBMMs as measured by reporter plasmid delivery. Percent lysis was determined by normalizing to Holin-Lysin and uninfected controls.

(C) Lysis of *L. monocytogenes* strains grown in BHI.

(D) Growth of cΔ*dacA* *L. monocytogenes* in WT (closed symbols) and Caspase 1^{-/-} (open symbols) BMMs.

Data are mean ± SEM and are representative of at least two independent experiments. *p < 0.05 (Students t test, two-tailed); NS, not significant.

To further characterize the effects of metabolic imbalance associated with c-di-AMP deficiency, a murine model of acute listeriosis was characterized. As previously reported (Witte et al., 2013), the cΔ*dacA* strain exhibited 3.8- and 4.6-log defects in both the spleen and liver, respectively (Figure 6G). When *citZ* was disrupted in the cΔ*dacA* strain, 2.7- and 2.4-logs of growth were rescued in the spleen and liver, respectively. Together with ex vivo analysis of intracellular growth, these findings support a model whereby metabolic imbalance associated with altered c-di-AMP levels specifically affects the magnitude of *L. monocytogenes* growth within the host cell cytosol with significant consequences on bacterial virulence.

Metabolic Imbalance in *L. monocytogenes* Causes Bacteriolysis within the Host Cytosol

To further elucidate the basis for the intracellular growth defect resulting from altered bacterial metabolic activity, we characterized the host response to infection. DNA released by *L. monocytogenes* leads to activation of Caspase 1 through the AIM2 inflammasome, resulting in pyroptosis during *L. monocytogenes* infection (Sauer et al., 2010). We found that bacteria with diminished levels of c-di-AMP production induce increased cell death during infection (Witte et al., 2013), while those with corrected metabolic activity through *citZ* disruption exhibited significantly decreased levels of host cell death (Fig-

ure 7A). Consistent with these observations, bacteria deficient in c-di-AMP production exhibit increased DNA delivery during infection, while the metabolic correction of this strain results in significantly reduced intracellular bacteriolysis (Figure 7B). Interestingly, no significant change in bacteriolysis is observed during growth in broth media (Figure 7C).

Pyroptosis is a protective host response to *L. monocytogenes* infection (Sauer et al., 2011). As such, we predicted that removal of this inflammatory response might rescue the growth of c-di-AMP deficient bacteria that exhibit c-di-AMP-related metabolic imbalance. Consistent with this hypothesis, intracellular growth of bacteria deficient in c-di-AMP production significantly improved in the absence of pyroptosis in primary bone-marrow-derived macrophages (Figure 7D). These observations suggest that the metabolic imbalance associated with diminished c-di-AMP signaling leads in part to diminished bacterial growth as a consequence of an enhanced restrictive host response. Together with our biochemical studies we conclude that these effects are most likely due to c-di-AMP-mediated regulation of LmPC and perhaps other metabolic factors.

DISCUSSION

Chemical proteomics provides a powerful approach to interrogate small molecule-protein interactions (Rix and Superti-Furga, 2009). The nucleotide interactome revealed here substantially broadens the known c-di-AMP protein-based signaling repertoire; identifying five previously unrecognized c-di-AMP-interacting proteins in *L. monocytogenes*, in addition to LmPC. The large number of effectors, their domain diversity, and seemingly disparate functional roles (Grinberg et al., 2006; Leigh and Dodsworth, 2007; Liu et al., 2006; Scott et al., 2004) are consistent with the pleiotropic effects of the *dacA* gene in *L. monocytogenes*. Our studies genetically segregate the effects of c-di-AMP on the magnitude of intracellular growth and intracellular bacterial lysis from other known phenotypes associated with loss of c-di-AMP production. We anticipate that the identification of several other protein receptors for c-di-AMP will guide future studies to define the molecular details governing changes in other c-di-AMP related phenotypes, including bacterial cell stability in broth, susceptibility to cell-wall targeting antibiotics, and bacterial growth rate in and out of the host.

Beyond broadening the known c-di-AMP effectors, our studies provide a detailed molecular characterization of the effects of c-di-AMP on PC activity. Our structural studies of apo and c-di-AMP-bound LmPC in comparison to known PC structures provide mechanistic insight into an unexpected mode of enzyme regulation. The nearly identical structure of the four LmPC monomers in the c-di-AMP complex is rather unusual among known PC structures. For SaPC, the differences in the orientations of the BC domains vary between 6° and 18° after their CT domains are superimposed (Xiang and Tong, 2008; Yu et al., 2009), and the differences are even larger for RePC (St Maurice et al., 2007). The structural differences among the four monomers in apo LmPC, as well as in SaPC and RePC, indicate that domain movements are necessary during PC catalysis, which is supported by recent EM studies on SaPC (Lasso et al., 2014). Based on these observations, we

favor a mechanism whereby c-di-AMP “freezes” the enzyme into a single (symmetrical) conformation, which is incompatible with catalysis.

Structure-function studies also clearly establish the location and strict specificity of the c-di-AMP-binding pocket. However, it is currently not clear how nucleotide specificity is achieved, as there are no hydrogen-bonding interactions to the base. Mutations of the sole base-interacting amino acid, Tyr722, do not extend inhibition activity to related nucleotides, and therefore other residues may contribute to this specificity. Beyond conferring specificity, changes in this pocket have significant consequences on enzyme function, even in the absence of ligand binding. These observations may be indicative of a previously unrecognized allosteric site for PC enzymes in response to c-di-AMP and possibly other stimuli. Significant variability is observed within the regulatory site identified by our studies. The EfPC-c-di-AMP interaction illustrates this mechanism of regulation extends beyond *L. monocytogenes* and that in the absence of amino acid conservation, divergent amino acids within the c-di-AMP-binding site can compensate to maintain nucleotide-mediated enzyme regulation. Given the central function of PC, proper integration of this enzyme into existing metabolic networks is required to maintain metabolic balance. As such, we postulate that the diversity of the sequence at this novel functional site may reflect divergent mechanisms of regulation, either through allosteric ligands or altered enzyme dynamics, which fine tune activity such that PC can properly integrate into the unique metabolic capacity of the diverse organisms that utilize this enzyme.

Bacteria have the capacity to thrive within a variety of environmental and host niches. This adaptability is largely reliant on the capacity to utilize diverse nutrients encountered in these disparate environments. The phosphoenolpyruvate-pyruvate-oxaloacetate node of central metabolism shapes the distribution of carbon metabolites among anabolic, catabolic, and energy-producing pathways (Jitrapakdee et al., 2008; Jitrapakdee and Wallace, 1999; Sauer and Eikmanns, 2005; Schär et al., 2010). Given this central function, aberrant metabolic activity as a consequence of altered c-di-AMP signaling may disrupt any one or a combination of the pathways required for bacterial adaptation to accessible nutrients. During infection, intracellular *L. monocytogenes* undergo a major metabolic transition from utilization of glucose to alternative carbon sources (Joseph et al., 2006), a process that affects activation of the master virulence regulator PrfA (Joseph et al., 2008; Stoll et al., 2008). Consequently, changes in PEP levels through increased pyruvate consumption may diminish PrfA activation and directly link c-di-AMP-related metabolic activity to virulence capacity. However, our observation that bacterial lysis is a downstream consequence of altered c-di-AMP-dependent metabolism suggests that the answer partially relates to the integrity of the bacterial cell. The metabolic burden generated by unregulated flux through the citric acid cycle may cause insufficiency in cell wall biosynthesis intermediates, leading to bacteriolysis. No apparent deficiency in de novo Asp, Ala, or Glx synthesis, each of which contributes to biosynthesis of the glycan and peptide components of the cell wall, suggests that limitations in these metabolites are not responsible. As the critical compound

utilized by *L. monocytogenes* to combat acid stress and the primary counter ion to osmotic stabilizing potassium (Booth and Higgins, 1990; Cotter et al., 2005; Yan et al., 1996), synthesis of glutamate must be tightly regulated to appropriately respond to changing conditions. As such, increased production of glutamate and the precursor metabolites in its biosynthetic pathway may have toxic effects on bacterial stability in the cytosol. Finally, hyperactivity of LmPC and subsequent turnover of OA by CitZ could lead to defects in lipid synthesis by consuming acetyl-CoA, a major precursor for the acyl-chain required to generate branched chain fatty acids. It is not known if *L. monocytogenes* strains deficient in these lipids exhibit enhanced intracellular lysis; however, the synthesis of these membrane components is required for *L. monocytogenes* survival within the cytosol of the host cell (Sun and O’Riordan, 2010). Clearly the effects of PC regulation extend beyond this intracellular pathogen. The evidence supporting PC regulation among lactic acid bacteria suggests that both extracellular pathogenic (i.e., *E. faecalis*) as well as nonpathogenic commensal *Lactobacilli* spp. utilize a similar mechanism of metabolic regulation, though the effects of this regulation remain an open question. Future studies to detail the functional consequences of c-di-AMP-mediated metabolic regulation through PC and likely other protein receptors will provide important insight into the physiology of these medically important host associated organisms.

In summary, the c-di-AMP-binding proteins reported here are widespread and conserved among environmental and pathogenic microbes alike and broadly expand the protein targets of c-di-AMP in bacteria. Furthermore, our biochemical and structural studies may provide important insight into a new molecular mechanism of central metabolic regulation through the enzyme PC. Finally, our studies demonstrate that c-di-AMP signaling regulates bacterial central metabolism and survival within the host environment. These findings lead us to propose that c-di-AMP serves a conserved and central role in the regulation of fundamental metabolic processes with important physiological implications in the unique environments occupied by the bacteria that utilize this secondary signaling molecule. A detailed understanding of bacterial signaling and metabolic regulation has profound practical and therapeutic applications. As a central element of these processes, the possibility of targeting c-di-AMP signaling pathways in this regard is an attractive option that remains to be explored.

EXPERIMENTAL PROCEDURES

C-di-AMP Pull-Downs

Soluble lysates from *L. monocytogenes* grown in 1.5 L BHI broth at 37°C to midexponential phase ($OD_{600} = 0.8$) were prepared in 30 ml of buffer (0.1 M Tris-HCl [pH 7.5], 150 mM NaCl, 0.1% v/v Tween-20, 1 mM PMSF). Lysates were incubated with rotation for 2 hr at 4°C with 100 μ l ethanoamine or c-di-AMP conjugated beads. Beads were washed three times with 5 ml PBS, mixed with 100 μ l of SDS-PAGE sample loading buffer, incubated at 56°C for 10 min, and the soluble fraction removed. The elution was repeated and each was pooled and processed for quantitative mass spectrometry as detailed in the Supplemental Information. Pull-downs were performed three independent times, one with the $\Delta pdeA$ strain and twice with WT *L. monocytogenes*.

Nucleotide-Binding Assays

The DRaCALA assays were performed with recombinant proteins or *E. coli* lysates in binding buffer (40 mM Tris, 100 mM NaCl, 20 mM MgCl₂ [pH 7.5]) and analyzed as described previously (Roelofs et al., 2011). For competition assays, 400 μM unlabeled nucleotide (ATP, GTP, cAMP, cGMP, NAD, NADH, c-di-AMP, c-di-GMP, pApA) was added prior to the addition of ³²P-c-di-AMP.

Protein Crystallization

LmPC crystals were grown by the sitting-drop vapor diffusion method at 20°C. For the c-di-AMP complex, the protein was incubated with 2.5 mM c-di-AMP and 2.5 mM ATP for 30 min at 4°C before setup. The reservoir solution contained 19% (w/v) PEG3350 and 0.2 M ammonium citrate (pH 7.0). The crystals appeared within 1–2 weeks and grew to full size after a few additional days. The crystals were cryoprotected in the reservoir solution supplemented with 10% (v/v) PEG200 and were flash-frozen in liquid nitrogen for data collection at 100 K. For the apo structure, the protein at 5 mg/ml was incubated with 2.5 mM ATP and 2.5 mM pyruvate for 30 min at 4°C before setup. The reservoir solution contained 16% (w/v) PEG3350, 0.1 M Bis-Tris (pH 6.5), and 1% (w/v) tacsimate (pH 7.0, Hampton). The crystals appeared within 1 week and grew to full size within a few more days. They were cryoprotected in the reservoir solution supplemented with 15% (v/v) ethylene glycol and flash frozen in liquid nitrogen. Data collection and structure determination were as detailed in the [Extended Experimental Procedures](#).

¹³C Incorporation Studies

Bacterial strains were grown overnight in BHI at 37°C. The cells were sedimented and washed with PBS. BHI was supplemented with 2.5 mg/ml [U-¹³C₆] glucose and were subsequently inoculated with an aliquot of the cell suspensions to an OD₆₀₀ of 0.1. The cultures were grown at 37°C until OD₆₀₀ = 1.0. Sodium azide was then added to a final concentration of 10 mM. The cells were centrifuged and washed three times with water. Bacterial cells (approximately 20 mg washed wet pellet) were suspended in 0.5 ml of 6 M hydrochloric acid. The mixture was heated at 105°C for 24 hr. The hydrolysate was placed on a column of Dowex 50W × 8 (H⁺ form; 200 to 400 mesh; 5 by 10 mm) previously washed with 4 ml of 7% formic acid and water. After loading the acid hydrolysate the column was washed with 4 ml of water. Sample was then eluted with 4 ml of 1 N ammonium hydroxide. The samples were air-dried under vacuum, and the residue was dissolved in 30 μl of pyridine. A total of 30 μl of (tert-butyl-dimethyl-silyl)-methyl-trifluoroacetamide containing 1% tert-butyl-dimethyl-silylchloride (Sigma) was added. The mixture was kept at 90°C for 60 min. Derivatized amino acids were analyzed by GC/MS as detailed in the [Extended Experimental Procedures](#).

Tissue Culture and Infection Assays

Primary bone-marrow-derived macrophages were generated as previously described (Jones and Portnoy, 1994). Macrophage growth curves, LDH, and DNA release assays were performed in primary or J2 immortalized BMMs (Sauer et al., 2010), as indicated. Plaque assays we conducted in L2 mouse fibroblasts (Sun et al., 1990). Murine infections were performed as described previously (Witte et al., 2013). All protocols were reviewed and approved by the Institutional Animal Care and Use Committee at the University of Wisconsin, Madison and the University of Washington.

ACCESSION NUMBERS

The Research Collaboratory for Structural Bioinformatics Protein Data Bank accession numbers for the new structures reported in this paper are 4QSH, 4QSK, and 4QSL. The Peptide Atlas database accession numbers for the LC MS/MS proteomics sequencing files are PASS00528.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Extended Experimental Procedures, seven figures, and one table and can be found with this article online at <http://dx.doi.org/10.1016/j.cell.2014.07.046>.

AUTHOR CONTRIBUTIONS

Conceived and designed the experiments: J.J.W., L.T., K.S. and P.H.C. Performed the experiments: J.J.W., L.T., K.S., P.H.C., T.N.H., A.R.J., M.P., M.D., N.P.O, D.P., and J.D.S. Contributed reagent/materials/analysis tools: A.T.I., Y.A.G, and M.S. Wrote the manuscript: J.J.W., L.T., K.S. and P.H.C.

ACKNOWLEDGMENTS

We would like to thank Dan Portnoy (UC Berkeley) for generous support and reagents, Brad Cookson and Dan Stetson (University of Washington) for murine bone marrow, Jason Smith and Sam Miller (University of Washington) for instrument access, Jake McKinlay (University of Indiana) for ¹³C-enrichment protocols, Nicholas Olivarez for technical support, and Joseph Mougous (University of Washington) for feedback on the preparation of the manuscript. We thank Linda Yu for earlier studies on PC expression and purification; Neil Whalen, Rick Jackimowicz and Howard Robinson for access to the X29A beamline at the NSLS. The in-house instrument for X-ray diffraction screening was purchased with an NIH grant to LT (OD012208). P.H.C. was also supported by an NIH Medical Scientist Training Program (GM007367). This work was supported by NIH grants U54AI057141 (to J.J.W.) and R01DK067238 (to L.T.).

Received: March 20, 2014

Revised: June 9, 2014

Accepted: July 29, 2014

Published: September 11, 2014

REFERENCES

- Bai, Y., Yang, J., Zhou, X., Ding, X., Eisele, L.E., and Bai, G. (2012). *Mycobacterium tuberculosis* Rv3586 (DacA) is a diadenylate cyclase that converts ATP or ADP into c-di-AMP. *PLoS ONE* 7, e35206.
- Bai, Y., Yang, J., Eisele, L.E., Underwood, A.J., Koestler, B.J., Waters, C.M., Metzger, D.W., and Bai, G. (2013). Two DHH subfamily 1 proteins in *Streptococcus pneumoniae* possess cyclic di-AMP phosphodiesterase activity and affect bacterial growth and virulence. *J. Bacteriol.* 195, 5123–5132.
- Bai, Y., Yang, J., Zarrella, T.M., Zhang, Y., Metzger, D.W., and Bai, G. (2014). Cyclic di-AMP impairs potassium uptake mediated by a cyclic di-AMP binding protein in *Streptococcus pneumoniae*. *J. Bacteriol.* 196, 614–623.
- Barker, J.R., Koestler, B.J., Carpenter, V.K., Burdette, D.L., Waters, C.M., Vance, R.E., and Valdivia, R.H. (2013). STING-dependent recognition of cyclic di-AMP mediates type I interferon responses during *Chlamydia trachomatis* infection. *MBio* 4, e00018–e00013.
- Benach, J., Swaminathan, S.S., Tamayo, R., Handelman, S.K., Folta-Stogniew, E., Ramos, J.E., Forouhar, F., Neely, H., Seetharaman, J., Camilli, A., and Hunt, J.F. (2007). The structural basis of cyclic diguanylate signal transduction by PilZ domains. *EMBO J.* 26, 5153–5166.
- Bernsel, A., Viklund, H., Hennerdal, A., and Elofsson, A. (2009). TOPCONS: consensus prediction of membrane protein topology. *Nucleic Acids Res.* 37 (Web Server issue), W465–W468.
- Booth, I.R., and Higgins, C.F. (1990). Enteric bacteria and osmotic stress: intracellular potassium glutamate as a secondary signal of osmotic stress? *FEMS Microbiol. Rev.* 6, 239–246.
- Campos, S.S., Ibarra-Rodriguez, J.R., Barajas-Ornelas, R.C., Ramirez-Guadiana, F.H., Obregón-Herrera, A., Setlow, P., and Pedraza-Reyes, M. (2014). Interaction of apurinic/aprimidinic endonucleases Nfo and ExoA with the DNA integrity scanning protein DisA in the processing of oxidative DNA damage during *Bacillus subtilis* spore outgrowth. *J. Bacteriol.* 196, 568–578.
- Corrigan, R.M., Abbott, J.C., Burhenne, H., Kaefer, V., and Gründling, A. (2011). c-di-AMP is a new second messenger in *Staphylococcus aureus* with a role in controlling cell size and envelope stress. *PLoS Pathog.* 7, e1002217.
- Corrigan, R.M., Campeotto, I., Jeganathan, T., Roelofs, K.G., Lee, V.T., and Gründling, A. (2013). Systematic identification of conserved bacterial c-di-AMP receptor proteins. *Proc. Natl. Acad. Sci. USA* 110, 9084–9089.

- Cotter, P.D., Ryan, S., Gahan, C.G., and Hill, C. (2005). Presence of GadD1 glutamate decarboxylase in selected *Listeria monocytogenes* strains is associated with an ability to grow at low pH. *Appl. Environ. Microbiol.* **71**, 2832–2839.
- Grinberg, I., Shteinberg, T., Gorovitz, B., Aharonowitz, Y., Cohen, G., and Borovok, I. (2006). The *Streptomyces* NrdR transcriptional regulator is a Zn ribbon/ATP cone protein that binds to the promoter regions of class Ia and class II ribonucleotide reductase operons. *J. Bacteriol.* **188**, 7635–7644.
- Huang, Y.H., Liu, X.Y., Du, X.X., Jiang, Z.F., and Su, X.D. (2012). The structural basis for the sensing and binding of cyclic di-GMP by STING. *Nat. Struct. Mol. Biol.* **19**, 728–730.
- Jitrapakdee, S., and Wallace, J.C. (1999). Structure, function and regulation of pyruvate carboxylase. *Biochem. J.* **340**, 1–16.
- Jitrapakdee, S., St Maurice, M., Rayment, I., Cleland, W.W., Wallace, J.C., and Attwood, P.V. (2008). Structure, mechanism and regulation of pyruvate carboxylase. *Biochem. J.* **413**, 369–387.
- Jones, S., and Portnoy, D.A. (1994). Characterization of *Listeria monocytogenes* pathogenesis in a strain expressing perfringolysin O in place of listeriolysin O. *Infect. Immun.* **62**, 5608–5613.
- Joseph, B., Przybilla, K., Stühler, C., Schauer, K., Slaghuis, J., Fuchs, T.M., and Goebel, W. (2006). Identification of *Listeria monocytogenes* genes contributing to intracellular replication by expression profiling and mutant screening. *J. Bacteriol.* **188**, 556–568.
- Joseph, B., Mertins, S., Stoll, R., Schär, J., Umeha, K.R., Luo, Q., Müller-Altrock, S., and Goebel, W. (2008). Glycerol metabolism and PrfA activity in *Listeria monocytogenes*. *J. Bacteriol.* **190**, 5412–5430.
- Kalia, D., Merey, G., Nakayama, S., Zheng, Y., Zhou, J., Luo, Y., Guo, M., Roembke, B.T., and Sintim, H.O. (2013). Nucleotide, c-di-GMP, c-di-AMP, cGMP, cAMP, (p)ppGpp signaling in bacteria and implications in pathogenesis. *Chem. Soc. Rev.* **42**, 305–341.
- Kim, H.J., Mittal, M., and Sonenshein, A.L. (2006). CcpC-dependent regulation of citB and Imo0847 in *Listeria monocytogenes*. *J. Bacteriol.* **188**, 179–190.
- Krasteva, P.V., Fong, J.C., Shikuma, N.J., Beyhan, S., Navarro, M.V., Yildiz, F.H., and Sondermann, H. (2010). *Vibrio cholerae* VpsT regulates matrix production and motility by directly sensing cyclic di-GMP. *Science* **327**, 866–868.
- Lasso, G., Yu, L.P., Gil, D., Lázaro, M., Tong, L., and Valle, M. (2014). Functional conformations for pyruvate carboxylase during catalysis explored by cryoelectron microscopy. *Structure* **22**, 911–922.
- Leigh, J.A., and Dodsworth, J.A. (2007). Nitrogen regulation in bacteria and archaea. *Annu. Rev. Microbiol.* **61**, 349–377.
- Liu, S., Bayles, D.O., Mason, T.M., and Wilkinson, B.J. (2006). A cold-sensitive *Listeria monocytogenes* mutant has a transposon insertion in a gene encoding a putative membrane protein and shows altered (p)ppGpp levels. *Appl. Environ. Microbiol.* **72**, 3955–3959.
- Luo, Y., and Helmann, J.D. (2012). Analysis of the role of *Bacillus subtilis* $\sigma(M)$ in β -lactam resistance reveals an essential role for c-di-AMP in peptidoglycan homeostasis. *Mol. Microbiol.* **83**, 623–639.
- Mehne, F.M., Gunka, K., Eilers, H., Herzberg, C., Kaever, V., and Stülke, J. (2013). Cyclic di-AMP homeostasis in *Bacillus subtilis*: both lack and high level accumulation of the nucleotide are detrimental for cell growth. *J. Biol. Chem.* **288**, 2004–2017.
- Nelson, J.W., Sudarsan, N., Furukawa, K., Weinberg, Z., Wang, J.X., and Breaker, R.R. (2013). Riboswitches in eubacteria sense the second messenger c-di-AMP. *Nat. Chem. Biol.* **9**, 834–839.
- Oppenheimer-Shaanan, Y., Wexselblatt, E., Katzhendler, J., Yavin, E., and Ben-Yehuda, S. (2011). c-di-AMP reports DNA integrity during sporulation in *Bacillus subtilis*. *EMBO Rep.* **12**, 594–601.
- Ouyang, S., Song, X., Wang, Y., Ru, H., Shaw, N., Jiang, Y., Niu, F., Zhu, Y., Qiu, W., Parvatykar, K., et al. (2012). Structural analysis of the STING adaptor protein reveals a hydrophobic dimer interface and mode of cyclic di-GMP binding. *Immunity* **36**, 1073–1086.
- Pozzi, C., Waters, E.M., Rudkin, J.K., Schaeffer, C.R., Lohan, A.J., Tong, P., Loftus, B.J., Pier, G.B., Fey, P.D., Massey, R.C., and O’Gara, J.P. (2012). Methicillin resistance alters the biofilm phenotype and attenuates virulence in *Staphylococcus aureus* device-associated infections. *PLoS Pathog.* **8**, e1002626.
- Rao, F., See, R.Y., Zhang, D., Toh, D.C., Ji, Q., and Liang, Z.X. (2010). YybT is a signaling protein that contains a cyclic dinucleotide phosphodiesterase domain and a GGDEF domain with ATPase activity. *J. Biol. Chem.* **285**, 473–482.
- Rix, U., and Superti-Furga, G. (2009). Target profiling of small molecules by chemical proteomics. *Nat. Chem. Biol.* **5**, 616–624.
- Roelofs, K.G., Wang, J., Sintim, H.O., and Lee, V.T. (2011). Differential radial capillary action of ligand assay for high-throughput detection of protein-metabolite interactions. *Proc. Natl. Acad. Sci. USA* **108**, 15528–15533.
- Sauer, U., and Eikmanns, B.J. (2005). The PEP-pyruvate-oxaloacetate node as the switch point for carbon flux distribution in bacteria. *FEMS Microbiol. Rev.* **29**, 765–794.
- Sauer, J.D., Witte, C.E., Zemansky, J., Hanson, B., Lauer, P., and Portnoy, D.A. (2010). *Listeria monocytogenes* triggers AIM2-mediated pyroptosis upon infrequent bacteriolysis in the macrophage cytosol. *Cell Host Microbe* **7**, 412–419.
- Sauer, J.D., Pereyre, S., Archer, K.A., Burke, T.P., Hanson, B., Lauer, P., and Portnoy, D.A. (2011). *Listeria monocytogenes* engineered to activate the NlrC4 inflammasome are severely attenuated and are poor inducers of protective immunity. *Proc. Natl. Acad. Sci. USA* **108**, 12419–12424.
- Schär, J., Stoll, R., Schauer, K., Loeffler, D.I., Eylert, E., Joseph, B., Eisenreich, W., Fuchs, T.M., and Goebel, W. (2010). Pyruvate carboxylase plays a crucial role in carbon metabolism of extra- and intracellularly replicating *Listeria monocytogenes*. *J. Bacteriol.* **192**, 1774–1784.
- Scott, J.W., Hawley, S.A., Green, K.A., Anis, M., Stewart, G., Scullion, G.A., Norman, D.G., and Hardie, D.G. (2004). CBS domains form energy-sensing modules whose binding of adenosine ligands is disrupted by disease mutations. *J. Clin. Invest.* **113**, 274–284.
- Shang, G., Zhu, D., Li, N., Zhang, J., Zhu, C., Lu, D., Liu, C., Yu, Q., Zhao, Y., Xu, S., and Gu, L. (2012). Crystal structures of STING protein reveal basis for recognition of cyclic di-GMP. *Nat. Struct. Mol. Biol.* **19**, 725–727.
- Shu, C., Yi, G., Watts, T., Kao, C.C., and Li, P. (2012). Structure of STING bound to cyclic di-GMP reveals the mechanism of cyclic dinucleotide recognition by the immune system. *Nat. Struct. Mol. Biol.* **19**, 722–724.
- Smith, W.M., Pham, T.H., Lei, L., Dou, J., Soomro, A.H., Beatson, S.A., Dykes, G.A., and Turner, M.S. (2012). Heat resistance and salt hypersensitivity in *Lactococcus lactis* due to spontaneous mutation of Ilmg_1816 (gdpP) induced by high-temperature growth. *Appl. Environ. Microbiol.* **78**, 7753–7759.
- St Maurice, M., Reinhardt, L., Surinya, K.H., Attwood, P.V., Wallace, J.C., Cleland, W.W., and Rayment, I. (2007). Domain architecture of pyruvate carboxylase, a biotin-dependent multifunctional enzyme. *Science* **317**, 1076–1079.
- Stoll, R., Mertins, S., Joseph, B., Müller-Altrock, S., and Goebel, W. (2008). Modulation of PrfA activity in *Listeria monocytogenes* upon growth in different culture media. *Microbiology* **154**, 3856–3876.
- Sun, Y., and O’Riordan, M.X. (2010). Branched-chain fatty acids promote *Listeria monocytogenes* intracellular infection and virulence. *Infect. Immun.* **78**, 4667–4673.
- Sun, A.N., Camilli, A., and Portnoy, D.A. (1990). Isolation of *Listeria monocytogenes* small-plaque mutants defective for intracellular growth and cell-to-cell spread. *Infect. Immun.* **58**, 3770–3778.
- Sundriyal, A., Massa, C., Samoray, D., Zehender, F., Sharpe, T., Jenal, U., and Schirmer, T. (2014). Inherent regulation of EAL domain-catalyzed hydrolysis of second messenger cyclic di-GMP. *J. Biol. Chem.* **289**, 6978–6990.
- Witte, G., Hartung, S., Büttner, K., and Hopfner, K.P. (2008). Structural biochemistry of a bacterial checkpoint protein reveals diadenylate cyclase activity regulated by DNA recombination intermediates. *Mol. Cell* **30**, 167–178.
- Witte, C.E., Whiteley, A.T., Burke, T.P., Sauer, J.D., Portnoy, D.A., and Woodward, J.J. (2013). Cyclic di-AMP is critical for *Listeria monocytogenes* growth, cell wall homeostasis, and establishment of infection. *MBio* **4**, e00282–e13.

- Woodward, J.J., Iavarone, A.T., and Portnoy, D.A. (2010). c-di-AMP secreted by intracellular *Listeria monocytogenes* activates a host type I interferon response. *Science* *328*, 1703–1705.
- Xiang, S., and Tong, L. (2008). Crystal structures of human and *Staphylococcus aureus* pyruvate carboxylase and molecular insights into the carboxyl-transfer reaction. *Nat. Struct. Mol. Biol.* *15*, 295–302.
- Yan, D., Ikeda, T.P., Shauger, A.E., and Kustu, S. (1996). Glutamate is required to maintain the steady-state potassium pool in *Salmonella typhimurium*. *Proc. Natl. Acad. Sci. USA* *93*, 6527–6531.
- Yang, J., Bai, Y., Zhang, Y., Gabrielle, V.D., Jin, L., and Bai, G. (2014). Deletion of the cyclic di-AMP phosphodiesterase gene (*cnpB*) in *Mycobacterium tuberculosis* leads to reduced virulence in a mouse model of infection. *Mol. Microbiol.* *93*, 65–79.
- Yin, Q., Tian, Y., Kabaleeswaran, V., Jiang, X., Tu, D., Eck, M.J., Chen, Z.J., and Wu, H. (2012). Cyclic di-GMP sensing via the innate immune signaling protein STING. *Mol. Cell* *46*, 735–745.
- Yu, L.P., Xiang, S., Lasso, G., Gil, D., Valle, M., and Tong, L. (2009). A symmetrical tetramer for *S. aureus* pyruvate carboxylase in complex with coenzyme A. *Structure* *17*, 823–832.
- Zhang, L., and He, Z.G. (2013). Radiation-sensitive gene A (RadA) targets DisA, DNA integrity scanning protein A, to negatively affect cyclic Di-AMP synthesis activity in *Mycobacterium smegmatis*. *J. Biol. Chem.* *288*, 22426–22436.
- Zhang, L., Li, W., and He, Z.G. (2013). DarR, a TetR-like transcriptional factor, is a cyclic di-AMP-responsive repressor in *Mycobacterium smegmatis*. *J. Biol. Chem.* *288*, 3085–3096.